

**A**

Reg. No. :

--	--	--	--	--	--	--	--	--	--

**Question Paper Code: 94021**

B.E./B.Tech. DEGREE EXAMINATION, APRIL 2024

Fourth Semester

Civil Engineering

19UMA423 - Numerical Methods

(Common to Chemical Engineering)

(Regulations 2019)

Duration: Three hours

Maximum: 100 Marks

Answer ALL Questions

**PART A - (10 x 1 = 10 Marks)**

1. Iteration method converges if  $|g^1(x)|$  \_\_\_\_\_ CO1-U  
(a)  $>1$       (b)  $<1$       (c)  $=0$       (d)  $>0$
2. Newton's method is also called method of \_\_\_\_\_ CO1-U  
(a) tangents      (b) slope      (c) secants      (d) false
3. Lagrange's interpolation formula can be used for \_\_\_\_\_ interval CO2-U  
(a) equal      (b) unequal      (c) equal and unequal      (d) none of these
4. Newton's divided difference formula used only for CO2-U  
\_\_\_\_\_ intervals  
(a) equal      (b) unequal      (c) equal and unequal      (d) none of these
5. Truncation error in Trapezoidal rule is of the order \_\_\_\_\_. CO3- U  
(a)  $h^3$       (b)  $h^2$       (c)  $h^4$       (d) 0
6. The Simpson's one third rule is approximated by \_\_\_\_\_. CO3- U  
(a) parabola      (b) trapezoid      (c) hyperbola      (d) elliptic
7. Taylor Series method will be very useful to give some \_\_\_\_\_ values for RK, Milne's and Adam's methods CO4-U  
(a) initial      (b) final      (c) intermediate      (d) two

8. In Euler's method, if h is large then it gives \_\_\_\_\_ value CO4-U  
 (a) accurate (b) inaccurate (c) average (d) None of these
9. PDE of second order, if  $B^2 - 4AC < 0$  then CO6-U  
 (a) parabolic (b) elliptic (c) hyperbolic (d) Non homogeneous
10. The equation  $u_{xx} + u_{yy} = 0$  CO5-App  
 (a) elliptic (b) parabolic (c) hyperbolic (d) Non homogeneous

PART – B (5 x 2= 10Marks)

11. What do you mean by diagonally dominant? CO1-U
12. State Newtons divided difference formula CO2-U
13. Evaluate  $\int_1^2 \frac{dx}{1+x^2}$  with 2 equal intervals using trapezoidal rule CO3-App
14. Using Taylor's series method find  $y(0.1)$  given  $y' = 1 + y$  with  $y(0) = 1$  CO4-App
15. Classify  $u_{xx} - 2u_{xy} + u_{yy} = 0$  CO6-U

PART – C (5 x 16= 80Marks)

16. (a) (i) Solve the equation  $x^3 - 2x - 5 = 0$  using iteration method CO1-App (8)  
 (ii) Solve  $x + 3y + 3z = 16$ ,  $x + 4y + 3z = 18$ ,  $x + 3y + 4z = 19$  CO1- App (8)  
 using Gauss Elimination method

Or

- (b) (i) Using Power method find numerically largest Eigen value and CO1- App (8)  
 the corresponding Eigen vector of the matrix 
$$\begin{pmatrix} 25 & 1 & 2 \\ 1 & 3 & 0 \\ 2 & 0 & -4 \end{pmatrix}$$
  
 (ii) Solve  $20x + y - 2z = 17$ ;  $3x + 20y - z = -18$ ;  $2x - 3y + 20z = 25$  using CO1- App (8)  
 Gauss Seidal method .

17. (a) (i) Using Lagrange's interpolation formula calculate the profit in CO2-App (8)  
 the year 2000 from the following data :

year	1997	1999	2001	2002
Profit ( Rs.in lakhs)	43	65	159	248

- (ii) Using Newton's divided difference formula find  $f(8)$  for the CO2-App (8)  
 data

X	4	5	7	10	11	13
Y	48	100	294	900	1210	2028

Or

- (b) (i) Using Newton's forward interpolation formula find  $f(2)$  for the CO2 -App (8) following data

X	0	5	10	15
Y	14	379	1444	3584

- (ii) Given the following table, find  $f(1.5)$  using cubic spline function CO2 -App (8)

x	1	2	3
$f(x)$	-8	-1	18

18. (a) (i) Compute the first and second derivatives of  $y$  at  $x = 4$  from CO3-App (8)

x	0	1	2	3	4
y	1	2.718	7.381	20.086	54.598

- (ii) Evaluate  $\int_0^6 \frac{1}{1+x^2} dx$  with 6 equal intervals by CO3-App (8)

- (a) Trapezoidal rule  
 (b) Simpson's  $\frac{1}{3}$  rule

Or

- (b) (i) Evaluate  $\int_{-1}^1 \frac{x^2}{1+x^4} dx$  using three point Gaussian quadrature formula. CO3-App (8)

- (ii) Evaluate  $\int_0^1 \int_0^1 \frac{1}{1+x+y} dxdy$  by using Simpson's rule by taking h=k=0.5 CO3-App (8)

19. (a) (i) Using Taylor's series method find  $y(1.1)$  given  $y' = x + y$  CO4-App (8) with  $y(1) = 0$

- (ii) Given  $\frac{dy}{dx} = 1 + y^2$ ,  $y(0) = 0$ ,  $y(0.2) = 0.2027$ ,  $y(0.4) = 0.4228$ ,  $y(0.6) = 0.6841$  evaluate  $y(0.8)$  by Adams – Bash forth Method.

Or

- (b) (i) Using R-K method of fourth order, solve  $\frac{dy}{dx} = \frac{y^2 - x^2}{y^2 + x^2}$  with CO4-App (8)  $y(0) = 1$  at  $x = 0.2$

- (ii) Given  $\frac{dy}{dx} = x^3 + y$ ,  $y(0) = 2$ ,  $y(0.2) = 2.443$ ,  $y(0.4) = 2.99$ ,  $y(0.6) = 3.68$ . Find  $y(0.8)$  by Milne's Predictor & Corrector method. CO4-App (8)

20. (a) (i) Solve  $\frac{\partial^2 u}{\partial x^2} = \frac{\partial u}{\partial t}$ ,  $u(0,t) = 0$ ,  $u(1,t) = 0$ ,  $u(x,0) = \sin \pi x$ . CO5-App (8)

Take  $h = 0.2$  and find the values of  $u$  up to  $t = 0.1$  using Bender-Schmidt's difference equation

(ii) Using Crank-Nicholson's difference equation to solve CO5-App (8)

$\frac{\partial^2 u}{\partial x^2} = \frac{\partial u}{\partial t}$   $u(0,t) = 0$ ,  $u(1,t) = t$ ,  $u(x,0) = 0$ . compute  $u$  for one time step function with  $h=0.25$ .

Or

(b) Solve the Laplace equation  $u_{xx} + u_{yy} = 0$  at the nine mesh points of the square given below. The values of  $u$  at the boundary are specified in the figure CO5- App (16)

0	11.1	17	19.7	18.6
0	$u_1$	$u_2$	$u_3$	21.9
0	$u_4$	$u_5$	$u_6$	21.0
0	$u_7$	$u_8$	$u_9$	17.0
0	8.7	12.1	12.8	9.0