

A**Reg. No. :**

--	--	--	--	--	--	--	--	--	--

Question Paper Code: 94021**B.E./B.Tech. DEGREE EXAMINATION, MAY 2022**

Fourth Semester

Civil Engineering

19UMA423 - Numerical Methods

(Common to Chemical Engineering)

(Regulations 2019)

Duration: Three hours

Maximum: 100 Marks

Answer ALL Questions

PART A - (10 x 1 = 10 Marks)

1. Iteration method converges if $|g'(x)|$ _____ CO1-U
(a) >1 (b) <1 (c) $=0$ (d) >0
2. Newton's method is also called method of _____ CO1-U
(a) tangents (b) slope (c) secants (d) false
3. Lagrange's interpolation formula can be used for _____ interval CO2-U
(a) equal (b) unequal (c) equal and unequal (d) none of these
4. In Newton's forward formula, $n=.$ CO2-U
(a) $\frac{x - x_0}{h}$ (b) $\frac{x - x_1}{h}$ (c) $\frac{x - x_2}{h}$ (d) $\frac{x - x_n}{h}$
5. Truncation error in Trapezoidal rule is of the order _____. CO3- U
(a) h^3 (b) h^2 (c) h^4 (d) 0
6. The Simpson's one third rule is approximated by _____. CO3- U
(a) parabola (b) trapezoid (c) hyperbola (d) elliptic
7. Taylor Series method will be very useful to give some _____ values for RK, Milne's and Adam's methods CO4-U
(a) initial (b) final (c) intermediate (d) two

8. In Euler's method, if h is large then it gives _____ value CO4-U
 (a) accurate (b) inaccurate (c) average (d) None of these
9. PDE of second order, if $B^2 - 4AC < 0$ then CO6-U
 (a) parabolic (b) elliptic (c) hyperbolic (d) Non homogeneous
10. The equation $u_{xx} + u_{yy} = 0$ CO5-App
 (a) elliptic (b) parabolic (c) hyperbolic (d) Non homogeneous

PART – B (5 x 2= 10Marks)

11. State the principle used in Gauss Elimination Method CO1-U
12. State Newtons divided difference formula CO2-U
13. Evaluate $\int_1^2 \frac{dx}{1+x^2}$ with 2 equal intervals using trapezoidal rule CO3-App
14. Using Taylor's series method find $y(0.1)$ given $y' = 1 + y$ with $y(0) = 1$ CO4-App
15. Classify $u_{xx} - 2u_{xy} + u_{yy} = 0$ CO6-U

PART – C (5 x 16= 80Marks)

16. (a) (i) Solve for a positive root of $3x - \cos x - 1 = 0$ using Newton's Raphson method correct to 6 decimal places. CO1-App (8)
 (ii) Solve $x + 3y + 3z = 16$, $x + 4y + 3z = 18$, $x + 3y + 4z = 19$ using Gauss Jordan method CO1- App (8)

Or

- (b) (i) Using Power method find numerically largest Eigen value and the corresponding Eigen vector of the matrix CO1- App (8)

$$\begin{pmatrix} 25 & 1 & 2 \\ 1 & 3 & 0 \\ 2 & 0 & -4 \end{pmatrix}$$
- (ii) Solve $20x+y-2z = 17$; $3x+20y-z = -18$; $2x-3y+20z = 25$ using CO1- App (8)
 Gauss Seidal method .
17. (a) (i) Using Lagrange's interpolation formula find $f(3)$ for the CO2-App (8)
 following data

X	0	1	2	5
Y	2	3	12	147

(ii) Using Newton's divided difference formula find $f(8)$ for the CO2-App data (8)

X	4	5	7	10	11	13
Y	48	100	294	900	1210	2028

Or

(b) (i) Using Newton's forward interpolation formula find $f(5)$ for the CO2 -App following data (8)

X	4	6	8	10
Y	1	3	8	10

(ii) Given the following table, find $f(1.5)$ using cubic spline CO2 -App function (8)

x	1	2	3
f(x)	-8	-1	18

18. (a) (i) Compute the first and second derivatives of y at $x = 1$ from CO3-App (8)

x	1	2	3	4
y	1	8	27	64

(ii) Evaluate $\int_0^6 \frac{1}{1+x^2} dx$ with 6 equal intervals by CO3-App (8)

(a) Trapezoidal rule

(b) Simpson's $\frac{1}{3}$ rule

Or

(b) (i) Evaluate. $\int_{-2}^2 e^{\frac{-x}{2}}$ using two point Gaussian quadrature CO3-App (8) formula

(ii) Evaluate $\int_1^{1.4} \int_2^{2.4} \frac{1}{xy} dx dy$ using Trapezoidal Rule with CO3-App (8)

$h = 0.1$ & $k = 0.1$

19. (a) (i) Using Taylor's series method find $y(1.1)$ given $y' = x + y$ CO4-App (8)
with $y(1) = 0$

(ii) Given $\frac{dy}{dx} = 1 + y^2$, $y(0) = 0$, $y(0.2) = 0.2027$, $y(0.4) = 0.4228$, CO4-App (8)

$y(0.6) = 0.6841$ evaluate $y(0.8)$ by Adams – Bash forth Method.

Or

(b) (i) Using R-K method of fourth order, solve $\frac{dy}{dx} = \frac{y^2 - x^2}{y^2 + x^2}$ with CO4-App (8)

$y(0) = 1$ at $x = 0.2$

(ii) Given $\frac{dy}{dx} = x^3 + y$, $y(0) = 2$, $y(0.2) = 2.443$, $y(0.4) = 2.99$, CO4-App (8)

$y(0.6) = 3.68$. Find $y(0.8)$ by Milne's Predictor & Corrector method.

20. (a) (i) Solve $\frac{\partial^2 u}{\partial x^2} = 32 \frac{\partial u}{\partial t}$, $u(0,t) = 0$, $u(1,t) = t$, $u(x,0) = 0$. Take CO5-App (8)
 $h = 0.25$ and find the values of u up to $t = 1$ using Bender-Schmidt's difference equation

(ii) Using Crank-Nicholson's difference equation to solve CO5-App (8)

$\frac{\partial^2 u}{\partial x^2} = \frac{\partial u}{\partial t}$ $u(0,t) = 0$, $u(1,t) = t$, $u(x,0) = 0$. compute u for one time step function with $h=0.25$.

Or

(b) Solve $\nabla^2 u = -10(x^2 + y^2 + 10)$ over the square mesh with sides CO5- App (16)

$x = 0$, $x = 3$, $y = 0$, $y = 3$ with $u=0$ on the boundary and mesh length 1 unit.