		Reg. No. :								
	Ques	tion Paper	Cod	e: 3	402	2				
B.]	E. / B.Tec	h. DEGREE E	XAM	INA	TIO	N, D	EC 2	2021		
		Fourth S	Semes	ster						
		Civil En	gineer	ring						
	01UN	IA422 - NUMI	ERIC	AL N	ЛЕТ	HOI	DS			
	(*	Common to EF	EE, EI	E an	d IC	E)				
		(Regulat	ion 2()13)						

Duration: Three hours

Maximum: 100 Marks

Answer ALL Questions.

PART A - $(10 \times 2 = 20 \text{ Marks})$

- 1. When is the convergence of an iterative method for solving the equation f(x) = 0 said to be (i) linear (ii) quadratic.
- 2. State the condition of convergence of Newton-Rapson method.
- 3. Give two indirect methods to solve a system of linear equations.
- 4. What do you mean by 'diagonally dominant'?
- 5. Define interpolation.
- 6. State Lagrange's interpolation formula.
- 7. State trapezoidal rule to evaluate $\int_{1}^{x_n} f(x) dx$.
- 8. Using Trapezoidal rule, evaluate $\int_0^{\pi} sinx \, dx$ by dividing the range into 6 equal parts.
- 9. Write the normal equations for fitting a straight line by the method of least squares.
- 10. State the principle of least squares.

- 11. (a) (i) Find the positive real root of 3x cosx 1 = 0 using Newton-Rapshon method. (8)
 - (ii) Solve the equation $x^3 + x^2 1 = 0$ for the positive root by iteration method. (8)

Or

- (b) (i) Using the secant method find a real root of the equation $f(x) = xe^x - 1 = 0.$ (8)
 - (ii) Find the real positive root of 3x cosx 1 = 0 by Newton Raphson method correct to 6 decimal places. (8)
- 12. (a) Solve the following system of equation using Gaussian elimination method. 28x + 4y - z = 32, x + 3y + 10z = 24, 2x + 17y + 4z = 35. (16)

Or

(b) (i) Solve the following system of equations by Gauss Seidel iteration method. 20x + y - 2z = 17, 3x + 20y - z = -18, 2x - 3y + 20z = 25 (8)

(ii) Using Jacobi method, find the eigen values and eigen vectors of $A = \begin{bmatrix} 1 & 3 \\ 3 & 4 \end{bmatrix}$. (8)

13. (a) Using Newton's backward formula find f(7.5) from the following table: (16)

X	1	2	3	4	5	6	7	8
f(x)	1	8	27	64	125	216	343	512

Or

- (b) (i) Using Newton's divided difference formula, find u(3) given u(1)=-26, u(2)=12, u(4)=256, u(6)=844. (8)
 - (ii) Using Newton's forward interpolation formula, find the polynomial f(x) satisfying the following data and hence find y(5). (8)

Х	4	6	8	10
у	1	3	8	10

14. (a) (i) Find the first two derivatives of $y = (x)^{1/3}$ at x = 50 & x = 56 given the table below.

x :	50	51	52	53	54	55	56
у:	3.6840	3.7084	3.7325	3.7563	3.7798	3.8030	3.8259

(ii) Evaluate
$$\int_0^{\frac{\pi}{2}} \int_0^{\frac{\pi}{2}} \sin(x+y) dx dy$$
 by using Trapezoidal rule and Simpson's rule.
(8)

Or

(b) (i) Compute first and second derivative of f(3) for the following data using difference table (8)

Х	3.0	3.2	3.4	3.6	3.8	4.0
<i>f</i> (x)	-14	-10.032	-5.296	-0.256	-6.672	14

(ii) Evaluate
$$\int_{0}^{1/2} \frac{2xy}{(1+x^2)(1+y^2)} dxdy$$
 using Trapezoidal rule with $h=k=0.25$. (8)

15. (a) (i) Find the equation of the best fitting straight line to the following data by method of group averages: (8)

Х	0	5	10	15	20	25	30
у	10	14	19	25	31	36	39

(ii) Fit a curve of the form $y = ae^{-bx}$ for the following data by the method of moments.

(8)

(8)

X	0	2	4	6	8	10
у	65	58	52	47	42	37

Or

(b) (i) Find a straight line fit of the form y = a + bx, by the method of group averages for the following data: (8)

x	0	5	10	15	20	25
у	12	15	17	22	24	30

(ii) By the method of moments, fit a straight line to the data.

X	1	2	3	4
Y	1.7	1.8	2.3	3.2

(8)